

Remarks

Claims 1-10, 12, 14 and 15 are pending. Claim 1 has been amended to eliminate the alkoxy, hydroxy, alkanoyloxy groups from the Markush group for R₇ and R₉. Claims 1 and 14 have been amended to delete the divalent group being Q-interrupted-Q-C₄-C₁₂alkylene-Q. Claim 1 has also been amended to delete two embedded preferred embodiments. Claims 3 and 5 have been amended to delete an embedded preferred embodiment. Claim 14 has been amended to restrict R₄ to groups that are bonded to the benzofuranone moiety by an oxygen atom. No new matter has been added.


The Examiner rejects claims 1-11 and 13-16 under 35 U.S.C. 112(2) as being indefinite for failing to particularly point out and distinctly claim the subject matter that Applicants regard as their invention. The Examiner objects to the phrase L is "Q interrupted...Q" in claims 1 and 14. The phrase has been deleted from both claims.

The objects/rejects claims 11, 13 and 16 as being "use claims". The claims have been cancelled without prejudice to the filing of divisional applications employing conventional process claim format.

The Examiner objects to embedded preferred embodiments in claims 1, 3, and 5. Each of the preferred embodiments has been deleted.

The Examiner rejects claims 1-5 and 7-11 (now claims 1-5 and 7-10) under 35 U.S.C. 102(b) as being anticipated by U.S. Pat. No. 5,981,160 ("Odenwalder et al."). Applicants respectfully traverse this rejection.

Odenwalder teaches the use of Benzofuranones of the formula:



as scavengers for the oxidized developer. The compounds shown in Odenwalder must contain an alkoxy group (OR₁) ortho to the point of attachment of the unfused phenyl moiety.

Claim 1 has been amended to exclude C₁-C₂alkoxy or C₂-C₂₀alkoxy (that can be interrupted) from the scope of R₇ and R₉. The R₇ and R₉ substituents are found on the substituted phenyl ring of formula II. In view of said amendment, Applicants request that the Examiner reconsider and withdraw her anticipation rejection of claims 1-5, 7-10 in view of Odenwalder.

The Examiner rejects claims 1-16 (now claims 1-10, 12, 14 and 15) under 35 U.S.C. 102 as being anticipated by U.S. Pat. No. 5,597,854 ("Birbaum et al."). Applicants respectfully traverse this rejection.

The present invention relates to a process for preventing migration of oxidized developer in a color photographic material by using the particularly defined benzofuranone compounds of claim 1 and its dependent claims. Birbaum describes the benzofuranone compounds as being possible co-additives with the latent light stabilizers taught therein. In fact, Birbaum listed these compounds as one of many optional types of UV absorbers and light stabilizers. The Examiner has not indicated any motivation for one skilled in the art to select the particularly defined compounds from the multiple of possible stabilizers for use in color photography films. The preferred applications don't even point to photography application. One skilled in the art would be expected to use the compositions taught in Birbaum for stabilizing plastics, rubbers, paints or adhesives. Applicants submit that the Examiner has selected portions of the prior art based upon an understanding of the claimed invention. The motivation and suggestion for use to prevent migration of oxidized developer in a color

photographic material is not found in Birbaum. In the absence of any such motivation, Birbaum certainly fails to disclose each and every element of the claimed process. Applicants request that the Examiner reconsider and withdraw her anticipation rejection of claims 1-10, 12, 14 and 15 in view of Birbaum.

The Examiner rejects claims 14-16 (now claims 14 and 15) under 35 U.S.C. 103 as being unpatentable over Birbaum in view of U.S. Pat. 4,325,863 ("Hinsken et al."). The Examiner applies Birbaum as described above. The Examiner cites Hinsken as teachings benzofuranone compounds wherein the R_2 and R_4 groups are pentyl radicals. The Examiner alleges that it would have been obvious to replace the butyl group taught in Birbaum with the pentyl groups of Hinsken. Applicants respectfully traverse this rejection.

Claim 14 has been amended to limit the R_4 group bonded to the benzofuranone moiety via an oxygen molecule. These compounds are described in examples 13-16 (compounds 29-32). All of the compounds in Hinsken are bonded to the benzofuranone moiety via a carbon atom. Hence, even assuming there is any motivation for the combination and modification urged by the Examiner, the resulting combination does not render the invention of claims 14 and 15 unpatentable.

Applicants submit that the present application is now in condition for allowance. Applicants request that the Examiner contact the undersigned representative if minor amendments will further prosecution.

Respectfully submitted,




David R. Crichton
Attorney for Applicants
Reg. No. 37,300

Ciba Specialty
Chemicals Corporation
540 White Plains Road
P.O. Box 2005
Tarrytown, New York 10591-9005
Tel: (914) 785-7124
Fax: (914) 785-7102


APPENDIX:

1. (amended) Process for preventing migration of the oxidised developer in a colour photographic material from one colour sensitive layer to another by incorporating a compound of the formula I into said material



wherein, if $n = 1$,

R_1 is a cyclic residue selected from naphthyl, phenanthryl, anthryl, 5,6,7,8-tetrahydro-2-naphthyl, 5,6,7,8-tetrahydro-1-naphthyl, thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, dibenzofuryl, chromenyl, xanthenyl, phenoxathiinyl, pyrrolyl, imidazolyl, pyrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, indolyl, indazolyl, purinyl, quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxaliny, quinazoliny, cinnoliny, pteridinyl, carbazolyl, -carboliny, phenanthridinyl, acridinyl, perimidiny, phenanthroliny, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl, biphenyl, terphenyl, fluorenyl or phenoxazinyl, each of which is unsubstituted or substituted by C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, hydroxy, halogen, amino, C_1 - C_4 alkylamino, phenylamino or $di(C_1$ - C_4 -alkyl)amino; or R_1 is a radical of formula II



and, if $n = 2$,

R_1 is unsubstituted or C_1 - C_4 alkyl- or hydroxy-substituted phenylene or naphthylene; or $-R_{12}$ - X - R_{13} ;

R_2 , R_3 , R_4 and R_5 are each independently of one another hydrogen; chloro; hydroxy; C_1 - C_{25} -alkyl; C_7 - C_9 phenylalkyl; unsubstituted or C_1 - C_4 alkyl-substituted phenyl; unsubstituted or C_1 - C_4 alkyl-

substituted C₅-C₈cycloalkyl; C₁-C₁₈alkoxy; C₁-C₁₈alkylthio; C₁-C₄alkylamino; di(C₁-C₄-alkyl)amino; C₁-C₂₅alkanoyloxy; C₁-C₂₅alkanoylamino; C₃-C₂₅alkenoyloxy; C₃-C₂₅alkanoyloxy which is


interrupted by oxygen, sulphur or >N-R_{14} ; C₆-C₉cycloalkylcarbonyloxy; benzoyloxy or C₁-

C₁₂alkyl-substituted benzoyloxy; or R₂ and R₃, or R₃ and R₄, or R₄ and R₅, together with the linking carbon atoms, form a benzene ring;

or R₄ is -C_mH_{2m}-COR_{15'}, -O-(C_vH_{2v})-COR_{15'}, -O-(CH₂)_q-OR_{32'}, -OCH₂-CH(OH)-CH₂-R_{15'},

-OCH₂-CH(OH)-CH₂-OR_{32'} or -(CH₂)_qOH;

or, if R₃, R₅ and R₆ are hydrogen, R₄ is additionally a radical of formula III



wherein R₁ is as defined above for n = 1;

R₆ is hydrogen or, when R₄ is hydroxy, R₆ can also be C₁-C₂₅alkyl or C₃-C₂₅alkenyl;

R₇ and R₉ are each independently of one another hydrogen; halogen; C₁-C₂₅alkyl; C₂-C₂₅alkyl

which is interrupted by oxygen, sulphur or >N-R_{14} ; C₁-C₂₅alkylthio; C₃-C₂₅-alkenyl; C₃-

C₂₅alkenyloxy; C₃-C₂₅alkynyl; C₃-C₂₅alkynyloxy; C₇-C₉phenylalkyl; C₇-C₉phenylalkoxy;

unsubstituted or C₁-C₄alkyl-substituted phenyl; unsubstituted or C₁-C₄alkyl- substituted

phenoxy; unsubstituted or C₁-C₄alkyl-substituted C₅-C₆cycloalkyl; unsubstituted or C₁-C₄alkyl-

substituted C₅-C₈cycloalkoxy; C₁-C₄alkylamino; di(C₁-C₄alkyl)amino; C₁-C₂₅alkanoyl; C₃-


C₂₅alkanoyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C₁-C₂₅alkanoylamino; C₃-

C₂₅alkenoyl; C₃-C₂₅alkenoyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C₃-C₂₅.

alkenoyloxy; C₃-C₂₅alkenoyloxy which is interrupted by oxygen, sulphur or >N-R_{14} ; C₆-C₉.

cycloalkylcarbonyl; C₆-C₉cycloalkylcarbonyloxy; benzoyl or C₁-C₁₂alkyl-substituted benzoyl;

benzoyloxy or C₁-C₁₂alkyl-substituted benzoyloxy; $\text{—O—}\underset{\text{R}_{19}}{\overset{\text{R}_{18}}{\text{C}}}\text{—}\overset{\text{O}}{\parallel}\text{C—R}_{15}$ or



R₇, R₈, R₉, R₁₀ and R₁₁ are each independently of one another hydrogen; halogen; hydroxy; C₁-

C₂₅alkyl; C₂-C₂₅alkyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C₁-C₂₅alkoxy; C₂-

C₂₅alkoxy which is interrupted by oxygen, sulphur or >N-R_{14} ; C₁-C₂₅alkylthio; C₃-C₂₅-alke-

nyl; C₃-C₂₅alkenyloxy; C₃-C₂₅alkynyl; C₃-C₂₅alkynyloxy; C₇-C₉phenylalkyl; C₇-C₉phenylalkoxy; unsubstituted or C₁-C₄alkyl-substituted phenyl; unsubstituted or C₁-C₄alkyl-substituted phenoxy; unsubstituted or C₁-C₄alkyl-substituted C₅-C₈cycloalkyl; unsubstituted or C₁-C₄alkyl-substituted C₅-C₈cycloalkoxy; C₁-C₄alkylamino; di(C₁-C₄alkyl)amino; C₁-C₂₅alkanoyl; C₃-

C₂₅alkanoyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C₁-C₂₅alkanoyloxy; C₃-

C₂₅alkanoyloxy which is interrupted by oxygen, sulphur or >N-R_{14} ; C₁-C₂₅alkanoylamino;

C₃-C₂₅alkanoyl; C₃-C₂₅alkanoyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C₃-C₂₅.

alkenoyloxy; C₃-C₂₅alkenoyloxy which is interrupted by oxygen, sulphur or >N-R_{14} ; C₆-C₉.

cycloalkylcarbonyl; C₆-C₉cycloalkylcarbonyloxy; benzoyl or C₁-C₁₂alkyl-substituted benzoyl;

benzoyloxy or C₁-C₁₂alkyl-substituted benzoyloxy; $\text{—O—}\overset{\overset{\text{R}_{18}}{|}}{\underset{\underset{\text{R}_{19}}{|}}{\text{C}}}\text{—}\overset{\overset{\text{O}}{||}}{\text{C}}\text{—R}_{15}$ or

$\text{—O—}\overset{\overset{\text{R}_{20}}{|}}{\underset{\underset{\text{H}}{|}}{\text{C}}}\text{—}\overset{\overset{\text{R}_{21}}{|}}{\underset{\underset{\text{R}_{22}}{|}}{\text{C}}}\text{—O—R}_{23}$ or, in formula II, R₇ and R₈, or R₈ and R₁₁, together with the linking

carbon atoms, form a benzene ring;

R₁₂ and R₁₃ are each independently of the other unsubstituted or C₁-C₄alkyl-substituted phenylene or naphthylene;



R₁₄ is hydrogen or C₁-C₈alkyl;

R₁₅ and R'₁₅ independently are hydroxy; $\left[\text{—O}^- \frac{1}{r} \text{M}^{r+} \right]$; C₁-C₂₀alkoxy; C₃-C₂₀alkoxy

interrupted by O and/or substituted by a radical selected from OH, phenoxy, C₇-

C₁₅alkylphenoxy, C₇-C₁₅alkoxyphenoxy; or are C₅-C₁₂cycloalkoxy; C₇-C₁₇phenylalkoxy; phenoxy;

$\text{—N}\begin{matrix} \nearrow \text{R}_{24} \\ \searrow \text{R}_{25} \end{matrix}$; or a group of the formula IIIa or IIIb



R₁₆ and R₁₇ are each independently of the other hydrogen, CF₃, C₁-C₁₂alkyl or phenyl, or R₁₆ and R₁₇, together with the linking carbon atom, are a C₅-C₈cycloalkylidene ring which is unsubstituted or substituted by 1 to 3 C₁-C₄alkyl;

R_{18} and R_{19} are each independently of the other hydrogen, C_1 - C_4 alkyl or phenyl;

R_{20} is hydrogen or C_1 - C_4 alkyl;

R_{21} is hydrogen; unsubstituted or C_1 - C_4 alkyl-substituted phenyl; C_1 - C_{25} alkyl; C_2 - C_{25} alkyl which is

interrupted by oxygen, sulphur or >N-R_{14} ; C_7 - C_9 phenylalkyl which is unsubstituted or

substituted at the phenyl moiety by 1 to 3 C_1 - C_4 alkyl; C_7 - C_{25} phenylalkyl which is interrupted by

oxygen, sulphur or >N-R_{14} and which is unsubstituted or substituted at the phenyl moiety

by 1 to 3 C_1 - C_4 alkyl; or R_{20} and R_{21} , together with the linking carbon atoms, form a C_3 -


C_{12} cycloalkylene ring which is unsubstituted or substituted by 1 to 3 C_1 - C_4 alkyl;

R_{22} is hydrogen or C_1 - C_4 alkyl;

R_{23} is hydrogen; C_1 - C_{25} alkanoyl; C_3 - C_{25} alkenoyl; C_3 - C_{25} alkanoyl which is interrupted by oxygen,

sulphur or >N-R_{14} ; C_2 - C_{25} alkanoyl which is substituted by a di(C_1 - C_6 alkyl)phosphonate


group; C_6 - C_9 cycloalkylcarbonyl; thenoyl; furoyl; benzoyl or C_1 - C_{12} alkyl-substituted benzoyl;





R_{24} and R_{25} are each independently of the other hydrogen or C_1 - C_{18} alkyl;

R_{26} is hydrogen or C_1 - C_8 alkyl;

R_{27} is a direct bond; C_1 - C_{18} alkylene; C_2 - C_{18} alkylene which is interrupted by oxygen, sulphur or



C_8 bicycloalkylene; unsubstituted or C_1 - C_4 alkyl-substituted phenylene;  or



R_{28} is hydroxy, $\left[-O^- \frac{1}{r} M^{r+} \right]$, C_1 - C_{18} alkoxy or $-N \begin{matrix} R_{24} \\ R_{25} \end{matrix}$;

R_{29} is oxygen or -NH-;

R_{30} is C_1 - C_{18} alkyl or phenyl;

R_{31} is hydrogen or C_1 - C_{18} alkyl;

R_{32} is C_1 - C_{18} alkanoyl; C_1 - C_8 alkanoyl substituted by phenyl or C_7 - C_{15} alkylphenyl; C_3 - C_{18} alkenoyl; cyclohexylcarbonyl; or naphthylcarbonyl;

L is a linking group of valency (k+1) and is as a divalent group

-O-;

Q- C_2 - C_{12} alkylene-Q;

-O-CH₂-CH(OH)-CH₂-O-;

-Q- C_2 - C_{12} alkylene-Q-CO-C_vH_{2v}-O-;

-O- C_2 - C_{12} alkylene-O-CH₂-CH(OH)-CH₂-O-;

~~Q-interrupted-Q- C_4 - C_{12} alkylene-Q-~~

Q-phenylene-Q or

Q-phenylene-D-phenylene-Q with D being C_1 - C_4 alkylene, O, S, SO or SO₂;

L as a trivalent group is Q-capped C_3 - C_{12} alkanetriyl, a trivalent residue of a hexose or a hexitol, or a group (-O-CH₂)₃C-CH₂OH; -Q-C₆H_{2a}-N(C₆H_{2b}-Q)-C₆H_{2c}-Q-;


-Q-C₃- C_{12} alkanetriyl(-Q-CO-C_vH_{2v}-O-)₂;

-O-C₃- C_{12} alkanetriyl(-O-CH₂-CH(OH)-CH₂-O-)₂; and

L as a tetravalent group is a tetravalent residue of a hexose or a hexitol;

-Q-C₄-C₁₂alkanetetryl(-Q-CO-C_vH_{2v}-O-)₃;

-O-C₄-C₁₂alkanetetryl(-O-CH₂-CH(OH)-CH₂-O-)₃; Q-capped C₄-C₁₂alkanetetryl; a group



M is an r-valent metal cation;


Q is oxygen or -NH-;

X is a direct bond, oxygen, sulphur or -NR₃₁-;

Z is a linking group of valency (k+1) and is as a divalent group C₂-C₁₂alkylene; Q-interrupted C₄-C₁₂alkylene; phenylene or phenylene-D-phenylene with D being C₁-C₄alkylene, O, S, SO or SO₂;

Z as a trivalent group is C₃-C₁₂alkanetriyl, a trivalent residue of a hexose or a hexitol, a group (-CH₂)₃C-CH₂OH, or a group -C_aH_{2a}-N(C_bH_{2b}-)-C_cH_{2c}-; and

Z as a tetravalent group is a tetravalent, carbon-ended residue of a hexose or a hexitol, C₄-



a, b, c and k independently are 1, 2 or 3;

m is 0 or a number from the range 1-12, preferably 1-6;

n is 1 or 2;

q is 1, 2, 3, 4, 5 or 6;

r is 1, 2 or 3; and


s is 0, 1 or 2;

v is 1, 2, 3, 4, 5, 6, 7 or 8, preferably 1 or 2;

provided that, when R₇ is hydroxy, alkanoyloxy or alkanoyloxy interrupted by O, S or N(R₁₄) and R₉ is hydrogen, R₁₀ is not identical with R₄; and when R₉ is hydroxy, alkanoyloxy or alkanoyloxy interrupted by O, S or N(R₁₄) and R₇ is hydrogen, R₈ is not identical with R₄.

3. (amended) Process according to claim 1 wherein in the compound of formula I

R_1 is naphthyl, phenanthryl, anthryl, 5,6,7,8-tetrahydro-2-naphthyl, 5,6,7,8-tetrahydro-1-naphthyl, thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, dibenzofuryl, chromenyl, xanthenyl, phenoxathiinyl, pyrrolyl, imidazolyl, pyrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, indolyl, indazolyl, purinyl, quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxaliny, quinazolinyl, cinnoliny, pteridinyl, carbazolyl, -carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl, biphenyl, terphenyl, fluorenyl or phenoxazinyl, each of which is unsubstituted or substituted by C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, hydroxy, halogen, amino, C_1 - C_4 alkylamino, phenylamino or di(C_1 - C_4 -alkyl)amino, or R_1 is a radical of formula II



and, if $n = 2$,


R_1 is unsubstituted or C_1 - C_4 alkyl- or hydroxy-substituted phenylene or naphthylene; or $-R_{12}-X-R_{13}-$,

R_2 , R_3 , R_4 and R_5 are each independently of one another hydrogen, chloro, hydroxy, C_1 - C_{25} -alkyl, C_7 - C_9 phenylalkyl, unsubstituted or C_1 - C_4 alkyl-substituted phenyl; unsubstituted or C_1 - C_4 alkyl-substituted C_5 - C_8 cycloalkyl; C_1 - C_{18} alkoxy, C_1 - C_{18} alkylthio, C_1 - C_4 alkylamino, di(C_1 - C_4 -alkyl)amino, C_1 - C_{25} alkanoyloxy, C_1 - C_{25} alkanoylamino, C_3 - C_{25} alkenoyloxy; C_3 - C_{25} alkanoyloxy which is

interrupted by oxygen, sulphur or $\text{>N}-R_{14}$; C_6 - C_9 cycloalkylcarbonyloxy, benzoyloxy or C_1 -

C_{12} alkyl-substituted benzoyloxy; or R_2 and R_3 , or R_3 and R_4 , or R_4 and R_5 , together with the linking carbon atoms, form a benzene ring; or R_4 is

$-C_mH_{2m}-COR_{15}$ or $-(CH_2)_qOH$ or, if R_3 , R_5 and R_6 are hydrogen, R_4 is additionally a radical of formula III



wherein R_1 is as defined above for $n = 1$;

R_6 is hydrogen or, when R_4 is hydroxy, R_6 can also be C_1 - C_{25} alkyl or C_3 - C_{25} alkenyl;

R_7 , R_8 , R_9 , R_{10} and R_{11} are each independently of one another hydrogen, halogen, hydroxy, C_1 -

C_{25} alkyl; C_2 - C_{25} alkyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C_1 - C_{25} alkoxy; C_2 -

C_{25} alkoxy which is interrupted by oxygen, sulphur or >N-R_{14} ; C_1 - C_{25} alkylthio, C_3 - C_{25} -alke-

nyl, C_3 - C_{25} alkenyloxy, C_3 - C_{25} alkynyl, C_3 - C_{25} alkynyloxy, C_7 - C_9 phenylalkyl, C_7 - C_9 phenylalkoxy, unsubstituted or C_1 - C_4 alkyl-substituted phenyl; unsubstituted or C_1 - C_4 alkyl- substituted phenoxy; unsubstituted or C_1 - C_4 alkyl-substituted C_5 - C_8 cycloalkyl; unsubstituted or C_1 - C_4 alkyl-substituted C_5 - C_8 cycloalkoxy; C_1 - C_4 alkylamino, di(C_1 - C_4 alkyl)amino, C_1 - C_{25} alkanoyl; C_3 -

C_{25} alkanoyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C_1 - C_{25} alkanoyloxy; C_3 -


C_{25} alkanoyloxy which is interrupted by oxygen, sulphur or >N-R_{14} ; C_1 - C_{25} alkanoylamino,

C_3 - C_{25} alkenoyl; C_3 - C_{25} alkenoyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C_3 - C_{25} -

alkenoyloxy; C_3 - C_{25} alkenoyloxy which is interrupted by oxygen, sulphur or >N-R_{14} ; C_6 - C_9 -

cycloalkylcarbonyl, C_6 - C_9 cycloalkylcarbonyloxy, benzoyl or C_1 - C_{12} alkyl-substituted benzoyl;

benzoyloxy or C₁-C₁₂alkyl-substituted benzoyloxy; $\text{---O---}\overset{\overset{\text{R}_{18}}{|}}{\underset{\underset{\text{R}_{19}}{|}}{\text{C}}}\text{---}\overset{\overset{\text{O}}{||}}{\text{C}}\text{---R}_{15}$ or




carbon atoms, form a benzene ring,

R₁₂ and R₁₃ are each independently of the other unsubstituted or C₁-C₄alkyl-substituted phenylene or naphthylene,

R₁₄ is hydrogen or C₁-C₈alkyl,

R₁₅ is hydroxy, $\left[\text{---O}^- \frac{1}{r} \text{M}^{r+} \right]$, C₁-C₂₀alkoxy, $\text{---N}\begin{matrix} \text{R}_{24} \\ \text{R}_{25} \end{matrix}$, or a group of the formula IIIa



R₁₆ and R₁₇ are each independently of the other hydrogen, CF₃, C₁-C₁₂alkyl or phenyl, or R₁₆ and R₁₇, together with the linking carbon atom, are a C₅-C₈cycloalkylidene ring which is unsubstituted or substituted by 1 to 3 C₁-C₄alkyl;

R₁₈ and R₁₉ are each independently of the other hydrogen, C₁-C₄alkyl or phenyl,

R₂₀ is hydrogen or C₁-C₄alkyl,


R₂₁ is hydrogen, unsubstituted or C₁-C₄alkyl-substituted phenyl; C₁-C₂₅alkyl; C₂-C₂₅alkyl which is

interrupted by oxygen, sulphur or >N---R_{14} ; C₇-C₉phenylalkyl which is unsubstituted or


substituted at the phenyl moiety by 1 to 3 C₁-C₄alkyl; C₇-C₂₅phenylalkyl which is interrupted by


oxygen, sulphur or >N---R_{14} and which is unsubstituted or substituted at the phenyl moiety

by 1 to 3 C₁-C₄alkyl, or R₂₀ and R₂₁, together with the linking carbon atoms, form a C₅-C₁₂cycloalkylene ring which is unsubstituted or substituted by 1 to 3 C₁-C₄alkyl;
R₂₂ is hydrogen or C₁-C₄alkyl,
R₂₃ is hydrogen, C₁-C₂₅alkanoyl, C₃-C₂₅alkenoyl; C₃-C₂₅alkanoyl which is interrupted by oxygen, sulphur or >N-R_{14} ; C₂-C₂₅alkanoyl which is substituted by a di(C₁-C₆alkyl)phosphonate group; C₆-C₉cycloalkylcarbonyl, thenoyl, furoyl, benzoyl or C₁-C₁₂alkyl-substituted benzoyl;



R₂₄ and R₂₅ are each independently of the other hydrogen or C₁-C₁₈alkyl,
R₂₆ is hydrogen or C₁-C₈alkyl,
R₂₇ is a direct bond, C₁-C₁₈alkylene; C₂-C₁₈alkylene which is interrupted by oxygen, sulphur or >N-R_{14} ; C₂-C₁₈alkenylene, C₂-C₂₀alkylidene, C₇-C₂₀phenylalkylidene, C₅-C₈cycloalkylene, C₇-

C₈bicycloalkylene, unsubstituted or C₁-C₄alkyl-substituted phenylene,  or



R_{28} is hydroxy, $\left[-O^- \frac{1}{r} M^{r+} \right]$, C_1 - C_{18} alkoxy or $-N \begin{matrix} R_{24} \\ R_{25} \end{matrix}$,

R_{29} is oxygen or -NH-,

R_{30} is C_1 - C_{18} alkyl or phenyl,

R_{31} is hydrogen or C_1 - C_{18} alkyl,

M is an r-valent metal cation,


Q is oxygen or -NH-,

X is a direct bond, oxygen, sulphur or $-NR_{31}-$,

Z is a linking group of valency (k+1) and is as a divalent group C_2 - C_{12} alkylene, Q-interrupted C_4 - C_{12} alkylene, phenylene or phenylene-D-phenylene with D being C_1 - C_4 alkylene, O, S, SO or SO_2 ;

Z as a trivalent group is C_3 - C_{12} alkanetriyl, a trivalent residue of a hexose or a hexitol, a group $(-CH_2)_3C-CH_2OH$, or a group $-C_aH_{2a}-N(C_bH_{2b})-C_cH_{2c}$; and

Z as a tetravalent group is a tetravalent residue of a hexose or a hexitol, C_4 - C_{12} alkanetetryl, a



a, b, c and k independently are 1, 2 or 3,

m is 0 or a number from the range 1-12, preferably 1-6,

n is 1 or 2,

q is 1, 2, 3, 4, 5 or 6,


r is 1, 2 or 3, and

s is 0, 1 or 2;

provided that, when R_7 is hydroxy, alkanoyloxy or alkanoyloxy interrupted by O, S or $N(R_{14})$ and

R_9 is hydrogen, R_{10} is not identical with R_4 .

5. (amended) Process according to claim 1 wherein the compound of formula I corresponds to the formula IV



wherein


R_2 is H or C_1 - C_{20} alkyl;

R_3 is H or C_1 - C_{18} alkyl;

R_4 is C_1 - C_8 alkyl, H, C_1 - C_6 alkoxy or a group $-C_mH_{2m}-COR_{15'}$; $-O-(C_vH_{2v})-COR_{15'}$;

$-O-(CH_2)_q-OR_{32'}$; $-OCH_2-CH(OH)-CH_2-R_{15'}$; $-OCH_2-CH(OH)-CH_2-OR_{32'}$;

or a group of the formula III



R_5 is H or C_1 - C_{18} alkyl;

R_7 and R_9 are each independently of one another hydrogen; halogen; C_1 - C_{25} alkyl; C_3 - C_{25} -alkenyl;


C_3 - C_{25} alkynyl; C_7 - C_9 phenylalkyl; unsubstituted or C_1 - C_4 alkyl-substituted phenyl; unsubstituted or

C_1 - C_4 alkyl-substituted C_5 - C_8 cycloalkyl;

R_8 , R_{10} and R_{11} independently are H, OH, chloro, C_1 - C_{18} alkyl, C_1 - C_{18} alkoxy, di(C_1 - C_4 alkyl)amino,



C_7 - C_9 phenylalkyl; unsubstituted or C_1 - C_4 alkyl-substituted phenyl; unsubstituted or C_1 - C_4 alkyl-

substituted C_5 - C_8 cycloalkyl; C_2 - C_{18} alkanoyloxy, C_3 - C_{18} -alkoxycarbonylalkoxy or




especially wherein at least 2 of the residues R_7 , R_8 , R_9 , R_{10} , R_{11} are H;

R_{15} is C_1 - C_{18} alkoxy; C_3 - C_{20} alkoxy interrupted by O; or are cyclohexyloxy; C_7 - C_{17} phenylalkoxy; phenoxy; or a group of formula IIIa or IIIb;



R_{16} and R_{17} , independently are H, C_1 - C_{12} alkyl or phenyl; or R_{16} and R_{17} , together with the bonding carbon atom form a C_5 - C_8 cycloalkylidene ring;

R_{20} , R_{21} and R_{22} independently are H or C_1 - C_4 alkyl;



R_{26} is C_1 - C_4 alkyl;

R_{32} is C_1 - C_{18} alkanoyl; C_1 - C_8 alkanoyl substituted by phenyl or C_7 - C_{15} alkylphenyl; C_3 - C_{18} alkenoyl; cyclohexylcarbonyl; or naphthylcarbonyl;

L is a divalent group -O-; $Q-C_2-C_{12}$ alkylene-Q; -O- CH_2 -CH(OH)- CH_2 -O-;

-Q- C_2 - C_{12} alkylene-Q-CO- C_vH_{2v} -O-; -O- C_2 - C_{12} alkylene-O- CH_2 -CH(OH)- CH_2 -O-;

Q is oxygen;

Z is C_2 - C_{12} alkylene;



k is 1;

m is 1, 2, 3, 4, 5 or 6;

v is 1 or 2; and

s is 0, 1 or 2.

14. (amended) Compound of the formula V or VI



wherein

~~R₄ is -(CH₂)₁₋₅, -COR'₁₅, or -CH(CH₃)-COR'₁₅, or -C_tH_{2t}-COR'₁₅, wherein C_tH_{2t} is a straight chain or branched alkylene moiety, or R₄ is -O-(C_vH_{2v})-COR'₁₅; -O-(CH₂)_q-OR'₃₂; -OCH₂-CH(OH)-CH₂-R'₁₅; or -OCH₂-CH(OH)-CH₂-OR'₃₂;~~



R'₇ is C₁-C₄alkyl and R'₈ is hydrogen or C₁-C₄alkyl;

R₁₅ is hydroxy, $\left[-O^- \frac{1}{r} M^{r+} \right]$, C₁-C₂₀alkoxy; C₃-C₂₀alkoxy interrupted by O and/or

substituted by a radical selected from OH, phenoxy, C₇-C₁₅alkylphenoxy, C₇-C₁₅alkoxyphenoxy;

or R₁₅ is C₅-C₁₂cycloalkoxy; C₇-C₁₇phenylalkoxy; phenoxy; $\begin{array}{c} R_{24} \\ \diagup \\ -N \\ \diagdown \\ R_{25} \end{array}$; or a group of formula

IIIa or IIIb;



~~R₁₅¹ is C₃-C₂₆ alkoxy interrupted by O and/or substituted by a radical selected from OH, phenoxy, C₇-C₁₅ alkylphenoxy, C₇-C₁₅ alkoxyphenoxy; or R₁₅¹ is C₅-C₁₂ cycloalkoxy, C₇-C₁₇ phenylalkoxy, phenoxy; or a group of formula IIIa or IIIb;~~

R₃₂ is C₁-C₁₈ alkanoyl; C₁-C₈ alkanoyl substituted by phenyl or C₇-C₁₅ alkylphenyl; C₃-C₁₈ alkenoyl; cyclohexylcarbonyl; or naphthylcarbonyl;

L is a linking group of valency (k+1) and is, as a divalent group,

-O-;

Q-C₂-C₁₂ alkylene-Q;

-O-CH₂-CH(OH)-CH₂-O-;

-Q-C₂-C₁₂ alkylene-Q-CO-C_vH_{2v}-O-;

-O-C₂-C₁₂ alkylene-O-CH₂-CH(OH)-CH₂-O-;

~~Q-interrupted Q-C₄-C₁₂ alkylene-Q;~~

Q-phenylene-Q or

Q-phenylene-D-phenylene-Q with D being C₁-C₄ alkylene, O, S, SO or SO₂;

L, as a trivalent group, is Q-capped C₃-C₁₂ alkanetriyl, a trivalent residue of a hexose or a hexitol, or a group (-O-CH₂)₃C-CH₂OH; -Q-C₆H_{2b}-N(C₆H_{2b}-Q)-C₆H_{2c}-Q-;


-Q-C₃-C₁₂ alkanetriyl(-Q-CO-C_vH_{2v}-O-)₂;

-O-C₃-C₁₂ alkanetriyl(-O-CH₂-CH(OH)-CH₂-O-)₂; and

L, as a tetravalent group, is a tetravalent residue of a hexose or a hexitol;

-Q-C₄-C₁₂ alkanetetriyl(-Q-CO-C_vH_{2v}-O-)₃;


-O-C₄-C₁₂alkanetetryl(-O-CH₂-CH(OH)-CH₂-O-)₃; Q-capped C₄-C₁₂alkanetetryl; a group



Q is oxygen or -NH-,

Z is a linking group of valency (k+1) and is as a divalent group C₂-C₁₂alkylene, Q-interrupted C₄-C₁₂alkylene, phenylene or phenylene-D-phenylene with D being C₁-C₄alkylene, O, S, SO or SO₂;
Z, as a trivalent group, is C₃-C₁₂alkanetriyl, a trivalent residue of a hexose or a hexitol, a group (-CH₂)₃C-CH₂OH, or a group -C_aH_{2a}-N(C_bH_{2b}-)-C_cH_{2c}-; and

Z, as a tetravalent group, is a tetravalent residue of a hexose or a hexitol, C₄-C₁₂alkanetetryl, a



a, b, c and k independently are 1, 2 or 3,

m is 0 or a number from the range 1-12,

s is 1 or 2,

~~and t is a number from the range 3-12,~~

v is 1, 2, 3, 4, 5, 6, 7 or 8;

and all other residues are as defined in claim 1 for formula I if n is 1.